

INFLUENCE OF BUBBLES ON NAVAL SYSTEMS OPERATING IN SHALLOW WATER: THE SCRIPPS PIER EXPERIMENT

Peter H. Dahl

Applied Physics Laboratory

College of Ocean and Fisheries Sciences

University of Washington

Seattle, Washington 98105

Telephone: (206) 543-2667

Fax: (206) 543-6785

E-Mail: dahl@apl.washington.edu

Award #: NAVSEA Contract N00039-96-D-0061, Task Order #2206002

Category of Research: high-frequency scattering, volume scattering

LONG-TERM GOALS

To predict the effects of bubbles, especially very high void fractions characteristic of surf zone regions, on MCM and weapons systems.

OBJECTIVES

The primary objective of the experiment was to measure properties of a bubble field, such as its spatial and temporal scales, its density, and its size distribution, within a near-shore environment. Data from this and future experiments will be interpreted along with the coastal oceanography and used to model the effect of bubbly environments on sonar system performance.

APPROACH

A collaborative, multi-institute experiment was performed in the vicinity of the Scripps Pier during Feb-March 1997, whose goal was the study of bubble field generation, transport, and distribution, as influenced by surf zone conditions. The collaboration involving NRL-SSC, IOS, APL-UW, SIO, NCPA, and NRL-DC, produced an ensemble of instrumentation at the site for measuring bubbles, ambient noise, temperature and salinity, currents and surface waves. APL-UW deployed an array of four upward-looking transducers (frequency 240kHz), which simultaneously measure vertical profiles of acoustic volume scattering from bubbles from four locations.

WORK COMPLETED

The Scripps Pier experiment and preliminary results of data analysis [1] were completed this year.

RESULTS

The transport of bubbles via rip currents emerged as a key feature surf zone bubble environment [1]. Images of volumetric backscattering strength vs time and depth (Fig. 1) reveal episodic events (by way of increased scattering level) which last $O(100)$ s and can occur in intervals of $O(10)$ min. Time lags for increased scattering at the four APL-UW locations were consistent with a seaward flow of rip currents equal to about 0.25 m/s.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Influence of Bubbles on Naval Systems Operating in Shallow Water: The Scripps Pier Experiment				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, 1013 N.E. 40th Street, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

IMPACT / APPLICATIONS

Data from the Scripps Pier Experiment are the most comprehensive to date that have been gathered in the surf zone environment. Results of this and perhaps future experiments this kind will no doubt have a strong influence on the design of sonar systems that must operate in very shallow water or surf zone environments.

TRANSITIONS

The results will be particularly useful to programs that address the influence of bubbles on naval systems operating near the sea surface, in the shallow water surf zone, or in vicinity of ship wakes.

RELATED PROJECTS

The work relates to other efforts within the ONR program entitled "Influence of Bubbles on Naval Systems Operating in Shallow Water", being carried out by NRL-SSC, IOS, and SIO-MPL.

REFERENCES

[1] P. H. Dahl, "Report on the Scripps Pier Bubble Experiment, Spring 1997," APL-UW TM 3-97, April 1997.

Fig.1 Depth vs time display of S_v (in decibels) as measured at 240kHz during coordinated Run 7 of the Scripps Pier Experiment. The plots are arranged such that their order, from top to bottom, represents the seaward direction, or increasing distance from shore. The abrupt increase in scattering level at minute 2.5 in the top plot is interpreted as a bubble cloud that subsequently spans the area delimited by the four transducer locations.

